

Contents lists available at [SciVerse ScienceDirect](http://SciVerse.ScienceDirect.com)

Applied Mathematics Letters

journal homepage: www.elsevier.com/locate/aml

Normal structure and moduli of UKK, NUC, and UKK* in Banach spaces

Satit Saejung^{a,b}, Ji Gao^{c,*}^a Department of Mathematics, Faculty of Science, Khon Kaen University, Khon Kaen 40002, Thailand^b Centre of Excellence in Mathematics, CHE, Sriyudthaya Road, Bangkok 10400, Thailand^c Department of Mathematics, Community College of Philadelphia, Philadelphia, PA 19130-3991, USA

ARTICLE INFO

Article history:

Received 15 September 2011

Received in revised form 6 January 2012

Accepted 9 January 2012

Keywords:

Modulus of UKK

Modulus of NUC

Modulus of UKK*

Normal structure

Uniform normal structure and ultraproduct space

ABSTRACT

Let X be a Banach space with the closed unit ball $B(X)$. In this paper, by directly extrapolating from the definitions of Uniformly Kadec–Klee (UKK), Nearly Uniformly Convex (NUC) and Weak* Uniformly Kadec–Klee (w^* UKK) spaces, we consider the concepts of the modulus of UKK and the modulus of NUC on X , and the modulus of UKK* on the dual space X^* of X . Some new properties of Banach spaces related to reflexivity and normal structure with the values of these moduli are obtained. Among these new results, we prove that if $B(X^*)$ is weak* sequentially compact and $\text{UKK}*((\frac{1}{\mu(X^*)})^-) > 1 - \frac{1}{\mu(X^*)}$ for X^* , then X has weak normal structure, where $\mu(X)$ is the separation measure of $B(X)$.

© 2012 Elsevier Ltd. All rights reserved.

1. Introduction

Let X be a normed linear space, and let $B(X) = \{x \in X : \|x\| \leq 1\}$, $S(X) = \{x \in X : \|x\| = 1\}$, and $B_\gamma(0) = \{x \in X : \|x\| < \gamma\}$ be the unit ball, the unit sphere, and open ball with radius γ of X respectively. Let X^* be the dual space of X .

Brodskiĭ and Mil'man introduced the concept of normal structure in 1948 [1].

For a reflexive Banach space, the normal structure and weak normal structure coincide.

In 1965, Kirk [2] proved that if a Banach space X has weak normal structure, then it has weak fixed point property, that is, every nonexpansive mapping from a weakly compact and convex subset of X into itself has a fixed point.

For a sequence $\{x_n\} \subseteq B(X)$, let $\text{co}(x_n)$, $\overline{\text{co}}(x_n)$, and $\overline{\text{co}}^w(x_n)$ be the convex hull, the closed convex hull, and the weak closed convex hull of the sequence $\{x_n\}$, respectively. Moreover, let $\text{sep}(x_n) \equiv \inf\{\|x_n - x_m\| : n \neq m\}$. We also write \xrightarrow{w} and $\xrightarrow{w^*}$ for the weak and the weak* convergence respectively, and write $f(a^-)$ for $\lim_{\varepsilon \rightarrow a^-} f(\varepsilon)$ for a function $f(\varepsilon)$.

Huff [3] introduced two concepts of Uniformly Kadec–Klee (UKK) spaces and Nearly Uniformly Convex (NUC) spaces on X in 1980.

Definition 1.1 ([3]). A Banach space X is called a UKK space (Uniform Kadec–Klee space) if for any $\varepsilon > 0$ there exists $0 < \delta < 1$ such that for any sequence $\{x_n\} \subseteq B(X)$ with $x_n \xrightarrow{w} x$ and $\text{sep}(x_n) \geq \varepsilon$, it follows that $x \in B_\delta(0)$.

Definition 1.2 ([3]). A Banach space X is called an NUC space (Nearly Uniform Convex space) if for any $\varepsilon > 0$ there exists $0 < \delta < 1$ such that for any sequence $\{x_n\} \subseteq B(X)$ with $\text{sep}(x_n) \geq \varepsilon$, it follows that $\text{co}(x_n) \cap B_\delta(0) \neq \emptyset$.

* Corresponding author. Tel.: +1 215 751 8688; fax: +1 215 751 8935.

E-mail addresses: saejung@kku.ac.edu (S. Saejung), jgao@ccp.edu (J. Gao).

He proved the relationship between UKK spaces and NUC spaces, and the relationship between NUC spaces and normal structure:

Theorem 1.3 ([3]). *Let X be a Banach space. Then X is an NUC space if and only if X is a UKK space and reflexive.*

Theorem 1.4 ([4,3]). *Every NUC space has normal structure.*

Dowling et al. [5] introduced the concept of UKK* spaces in 2008.

Definition 1.5 ([5]). Let X be a Banach space. The dual space X^* of X is called a UKK* space (Uniform Kadec–Klee* space) if for any $\varepsilon > 0$ there exists $0 < \delta < 1$ such that for any sequence $\{f_n\} \subseteq B(X^*)$ with $f_n \xrightarrow{w^*} f$ and $\text{sep}(f_n) \geq \varepsilon$, it follows that $f \in B_\delta(0)$ for X^* .

The following result was proved:

Theorem 1.6 ([5]). *Let X be a Banach space such that $B(X^*)$ is weak* sequentially compact. If X^* is a UKK* space, then X has weak fixed point property.*

Some measures of noncompactness on a Banach space X have been introduced to study the geometrical properties of Banach spaces, see [6–12].

Definition 1.7 ([6]). Let A be a bounded set of a metric space. The α -measure, χ -measure, and β -measure of noncompactness of A are defined by

$$\alpha(A) = \inf\{\varepsilon > 0 : A \text{ can be covered by finitely many sets with diameter } \leq \varepsilon\};$$

$$\chi(A) = \inf\{\varepsilon > 0 : A \text{ can be covered by finitely many balls with radii } \leq \varepsilon\};$$

$$\beta(A) = \sup\{\varepsilon > 0 : A \text{ contains an infinite } \varepsilon\text{-separation, that is, there is an infinite subset } C \text{ of } A \text{ such that } d(x, y) \geq \varepsilon \text{ for all } x, y \in C \text{ with } x \neq y\},$$

respectively.

It is known that if A is a bounded subset of a Banach space, then

$$\chi(A) \leq \beta(A) \leq \alpha(A) \leq 2\chi(A).$$

Definition 1.8. Let X be a Banach space. The α -modulus [6,10], χ -modulus [6,7], and β -modulus [6,9] of noncompact convexity of X are defined by

$$\Delta_\alpha(\varepsilon) = \inf\{1 - \inf_{x \in A} \|x\| : A \text{ is a convex subset of } B(X) \text{ and } \alpha(A) > \varepsilon\};$$

$$\Delta_\chi(\varepsilon) = \inf\{1 - \inf_{x \in A} \|x\| : A \text{ is a convex subset of } B(X) \text{ and } \chi(A) > \varepsilon\};$$

$$\Delta_\beta(\varepsilon) = \inf\{1 - \inf_{x \in A} \|x\| : A \text{ is a convex subset of } B(X) \text{ and } \beta(A) > \varepsilon\},$$

respectively.

The following result was proved:

Theorem 1.9 ([6, Theorem V.1.7]). *Let X be a Banach space. If $\Delta_\chi(1) > 0$, then X is reflexive.*

The function $P_X(\varepsilon)$ below was defined by Partington: (it was also considered in [12] but denoted by $\diamond(\varepsilon)$)

Definition 1.10 ([6,13]).

$$P_X(\varepsilon) = \inf\{1 - \|x\| : x_n \xrightarrow{w} x, \{x_n\} \subseteq B(X) \text{ and } \text{sep}(x_n) > \varepsilon\}.$$

The following result was proved:

Theorem 1.11 ([6, Theorem V.1.10, Remark V.1.12]). *For a reflexive Banach space X , $\Delta_\beta(\cdot) = P_X(\cdot) = \diamond(\cdot)$.*

2. Main results

Lemma 2.1 ([14]). *Let X be a Banach space. Suppose that X does not have weak normal structure. Then there exists a weak null sequence $\{y_n\} \subseteq X$ such that for every $k \in \mathbb{N}$,*

$$1 - \frac{1}{k} < \left\| \sum_{j=0}^l \lambda_j y_{k+j} - y_{k+l+1} \right\| < 1 + \frac{1}{k}$$

holds for all $l \in \mathbb{N}$ and all $\lambda_1, \dots, \lambda_l \geq 0$ with $\sum_{j=1}^l \lambda_j = 1$.

Then, it is easy to get the following result:

Lemma 2.2. Let X be a Banach space without weak normal structure. Then for any $0 < \epsilon < 1$, there exists a sequence $\{x_n\} \subseteq S(X)$ with $x_n \xrightarrow{w} 0$, and

$$1 - \epsilon < \|x_{n+1} - x\| < 1 + \epsilon$$

for sufficiently large n , and any $x \in \text{co}\{x_k\}_{k=1}^n$.

Theorem 2.3 ([15]). Let X be an infinite dimensional normed space and $\mu(X) = \beta(B(X))$. Then $1 \leq \mu(X) \leq 2$, and $\mu(l_p) = 2^{\frac{1}{p}}$ where $1 \leq p < \infty$.

By directly extrapolating from the definitions of UKK and NUC spaces, we first consider the concepts of modulus of UKK and modulus of NUC on X as follows:

Definition 2.4. Let X be a Banach space. For $0 < \epsilon \leq \mu(X)$, let

$$\text{UKK}(\epsilon) = 1 - \delta,$$

where δ is the smallest number in $(0, 1]$ such that $x \in B_\delta(0)$ whenever $\{x_n\} \subseteq B(X)$ satisfies $x_n \xrightarrow{w} x$ and $\text{sep}(x_n) \geq \epsilon$. For $\mu(X) < \epsilon \leq 2$, let $\text{UKK}(\epsilon) = 1$. Then the function $\text{UKK}(\epsilon)$ is called the UKK modulus of X .

Remark 2.5. It follows directly from the definitions that

$$\text{UKK}(\epsilon) = P_X(\epsilon) = \diamond(\epsilon)$$

for $0 \leq \epsilon \leq \mu(X)$.

Definition 2.6. Let X be a Banach space. For $0 < \epsilon \leq \mu(X)$, let

$$\text{NUC}(\epsilon) = 1 - \delta,$$

where δ is the smallest number in $(0, 1]$ such that $\text{co}(x_n) \cap B_\delta(0) \neq \emptyset$ whenever $\{x_n\} \subseteq B(X)$ satisfies $\text{sep}(x_n) \geq \epsilon$. For $\mu(X) < \epsilon \leq 2$, let $\text{NUC}(\epsilon) = 1$. Then the function $\text{NUC}(\epsilon)$ is called the NUC modulus of X .

We then consider the following concept for the dual space X^* :

Definition 2.7. Let X^* be a dual of a Banach space X . For $0 < \epsilon \leq \mu(X^*)$, let

$$\text{UKK}^*(\epsilon) = 1 - \delta,$$

where δ is the smallest number in $(0, 1]$ such that $f \in B_\delta(0)$ whenever $\{f_n\} \subseteq B(X^*)$ satisfies $f_n \xrightarrow{w^*} f$ and $\text{sep}(f_n) \geq \epsilon$. For $\mu(X^*) < \epsilon \leq 2$, let $\text{UKK}^*(\epsilon) = 1$. Then the function $\text{UKK}^*(\epsilon)$ is called the UKK^* modulus of X^* .

Example 2.8. (1) $\text{NUC}(2) = 0$ for l_1 space. Let $e_i = (0, \dots, 0, 1, 0, \dots)$, where the i th element is 1, be the standard basis of l_1 . Then $\text{sep}(e_i) = 2$, and $\text{co}(e_i) \subseteq S(l_1)$. So $\mu(l_1) = 2$ and $\text{NUC}(2) = 0$.

(2) $\text{NUC}(\epsilon) = 1 - \sqrt{1 - \frac{\epsilon^2}{4}}$ for Hilbert spaces.

We immediately obtain the following easy results.

Proposition 2.9. Let X be a Banach space.

- (1) All UKK modulus, NUC modulus of X , and UKK^* modulus of X^* are nondecreasing functions.
- (2) X is a UKK space if and only if $\text{UKK}(\epsilon) > 0$ for all $\epsilon > 0$.
- (3) X is an NUC space if and only if $\text{NUC}(\epsilon) > 0$ for all $\epsilon > 0$.
- (4) X^* is a UKK^* space if and only if $\text{UKK}^*(\epsilon) > 0$ for all $\epsilon > 0$.

The following ingenious result was proved by James.

Theorem 2.10 ([16]). Let X be a Banach space. Then X is not reflexive if and only if for any $0 < \epsilon < 1$, there are a sequence $\{x_n\} \subseteq S(X)$ and a sequence $\{f_n\} \subseteq S(X^*)$ such that

- (a) $\langle x_m, f_n \rangle = \epsilon$ whenever $n \leq m$; and
- (b) $\langle x_m, f_n \rangle = 0$ whenever $n > m$.

Using the preceding result, we obtain the following result.

Theorem 2.11. If X is a Banach space with $\text{NUC}(1^-) > 0$, then it is reflexive.

Proof. By the assumption, we can find an $\varepsilon \in (0, 1)$ such that $\text{NUC}(\varepsilon) > 1 - \varepsilon$. Suppose that X is not reflexive. By Theorem 2.10, there are two sequences $\{x_n\} \subseteq S(X)$ and $\{f_n\} \subseteq S(X^*)$ satisfying the two conditions there. Then $\|x_n - x_m\| \geq \langle x_n - x_m, f_n \rangle = \varepsilon$ for $n > m$ and hence $\text{sep}(x_n) = \inf\{\|x_n - x_m\| : n > m\} \geq \varepsilon$.

Let $\alpha_1, \alpha_2, \dots, \alpha_k \geq 0$ be such that $\sum_{i=1}^k \alpha_i = 1$. Suppose that k_0 is the first number such that $\alpha_{k_0} > 0$. Then $\|\sum_{i=1}^k \alpha_i x_i\| \geq \langle \sum_{i=1}^k \alpha_i x_i, f_{k_0} \rangle = \varepsilon$ and this implies that $\text{co}(x_n) \cap B_\varepsilon(0) = \emptyset$. So $\text{NUC}(\varepsilon) < 1 - \varepsilon$, a contradiction. \square

The following theorem shows the relation between the moduli $\text{UKK}(\varepsilon)$ and $\text{NUC}(\varepsilon)$:

Theorem 2.12. Let X be a Banach space. Then

1. $\text{NUC}(\varepsilon) \leq \text{UKK}(\varepsilon)$ for any $0 < \varepsilon < 1$;
2. $\text{NUC}(\varepsilon) = \text{UKK}(\varepsilon)$ for any $0 < \varepsilon < 1$ if X is reflexive;
3. For any $0 < \varepsilon < 1$, $\text{NUC}(\varepsilon) = \alpha > 0$ if and only if $\text{UKK}(\varepsilon) = \alpha > 0$ and X is reflexive.

Proof. (1) Put $\alpha \equiv \text{NUC}(\varepsilon)$. Let $\eta > 0$ and $\{x_n\} \subseteq S(X)$ be such that $\text{sep}(x_n) \geq \varepsilon$ and $x_n \xrightarrow{w} x$. If $\{x_{n_k}\}$ is a subsequence of $\{x_n\}$, then $\text{sep}(x_{n_k}) \geq \varepsilon$ and $x_{n_k} \xrightarrow{w} x$. We can find sequences $\{p_n\}$ and $\{q_n\}$ of positive integers, a sequence $\{\alpha_n\}$ in $[0, 1]$ such that the following conditions are satisfied:

- $1 < p_1 < q_1 < p_2 < q_2 < \dots < p_j < q_j < \dots$ for all j ;
- $\sum_{i=p_j}^{q_j} \alpha_i = 1$ for all j ;
- $\sum_{i=p_j}^{q_j} \alpha_i x_i \in B_{1-\alpha+\eta}(0)$ for all j .

It is easy to see that $\sum_{i=p_j}^{q_j} \alpha_i x_i \xrightarrow{w} x$ as $j \rightarrow \infty$. Hence $x \in B_{1-\alpha+\eta}(0)$ for any $\eta > 0$ and we have $\text{UKK}(\varepsilon) \geq \alpha$.

(2) Put $\beta \equiv \text{UKK}(\varepsilon)$. Suppose that a sequence $\{x_n\} \subseteq S(X)$ satisfies $\text{sep}(x_n) \geq \varepsilon$. By the reflexivity and passing to an appropriate subsequence but still denoted by $\{x_n\}$, we may assume that $x_n \xrightarrow{w} x \in \overline{\text{co}}^w(x_n) = \overline{\text{co}}(x_n)$. It follows that $x \in \text{co}(x_n) \cap B_{1-\beta+\eta}(0)$ for any $\eta > 0$. Since η can be arbitrarily small, we have $\text{NUC}(\varepsilon) \geq \beta$.

(3) This is a direct result of (2) and Theorem 2.11. \square

The following is the similar relationship between UKK spaces and NUC spaces:

Corollary 2.13 ([3]). Let X be a Banach space. Then X is an NUC space if and only if X is a UKK space and reflexive.

Proof. This is a direct result of Theorem 2.12. \square

The following result is closely related to the one in [17].

Theorem 2.14. Let X be a Banach space. If $\text{UKK}(1^-) > 0$, then X has weak normal structure.

Proof. By the assumption, there is $\eta \in (0, 1)$ such that $\text{UKK}(\frac{1-\eta}{1+\eta}) > \frac{\eta}{1+\eta}$. Suppose that X does not have weak normal structure. By Lemma 2.2, there is a weak null sequence $\{x_n\} \subseteq S(X)$ satisfying the condition:

$$1 - \eta < \|x_n - x_m\| < 1 + \eta \quad \text{for } n \neq m.$$

Consider the sequence

$$y_n \equiv \frac{x_1 - x_{n+1}}{1 + \eta}.$$

Then $\{y_n\} \subseteq B(X)$ and $\|y_m - y_n\| = \frac{\|x_{n+1} - x_{m+1}\|}{1 + \eta} > \frac{1-\eta}{1+\eta}$ for $m \neq n$. So $\text{sep}(y_n) \geq \frac{1-\eta}{1+\eta}$. It is easy to see that $y_n \xrightarrow{w} \frac{x_1}{1+\eta}$. This implies that $\text{UKK}(\frac{1-\eta}{1+\eta}) \leq \frac{\eta}{1+\eta}$, a contradiction. \square

Corollary 2.15 ([3]). If X is an NUC space, then X has normal structure.

Corollary 2.16 ([3]). If X is a UKK space, then X has weak normal structure.

In 2007, Saejung proved the following result:

Lemma 2.17 ([18]). If X is a Banach space with $B(X^*)$ weak* sequentially compact and it fails to have weak normal structure, then for any $\varepsilon > 0$ there are $\{x_1, x_2, \dots, x_n\} \subseteq S(X)$ and $\{f_1, f_2, \dots, f_n\} \subseteq S(X^*)$ such that

- (a) $\|x_i - x_j\| - 1 < \varepsilon$, for all $i \neq j$;
- (b) $\langle x_i, f_i \rangle = 1$, for all $1 \leq i \leq n$; and
- (c) $|\langle x_j, f_i \rangle| < \varepsilon$, for all $i \neq j$.

This result can be extended as follows:

Lemma 2.18. *If X is a Banach space with $B(X^*)$ weak* sequentially compact and it fails to have weak normal structure, then for any $\varepsilon > 0$ there are two sequences $\{x_n\} \subseteq S(X)$ and $\{f_n\} \subseteq S(X^*)$ such that*

- (a) $\|x_i - x_j\| - 1 < \varepsilon$, for all $i \neq j$;
- (b) $\langle x_i, f_i \rangle = 1$, for all $i \in \mathbb{N}$;
- (c) $|\langle x_j, f_i \rangle| < \varepsilon$, for all $i \neq j$; and
- (d) $\|f_i - f_j\| > 1 - \varepsilon$, for all $i \neq j$.

Proof. Let $\varepsilon > 0$, from Lemma 2.2 and the assumptions, for $\eta = \frac{\varepsilon}{2}$ we can find sequences $\{x_n\} \subseteq S(X)$, $\{f_n\} \subseteq S(X^*)$ and $f \in B(X^*)$, such that

- (a) $\|x_i - x_j\| - 1 < \eta$, where $i \neq j$;
- (b) $\langle x_i, f_i \rangle = \|x_i\| = 1$, where $1 \leq i < \infty$;
- (c) $x_n \xrightarrow{w} 0$ and
- (d) $f_n \xrightarrow{w^*} f$.

Since $x_n \xrightarrow{w} 0$, we can find a subsequence $1 < k_1 < k_2 < \dots < k_n < \dots$ such that $|\langle x_m, f_1 \rangle| < \eta$ for $m > k_1$, $|\langle x_m, f_{k_1} \rangle| < \eta$ for $m > k_2$, and so on.

So, the subsequence $1 < k_1 < k_2 < \dots < k_n < \dots$ satisfies $|\langle x_{k_j}, f_{k_i} \rangle| < \eta$ for $i < j$.

Without loss of generality, we still use \mathbb{N} to denote the subsequence $1 < k_1 < k_2 < \dots < k_n < \dots$. So we have $|\langle x_j, f_i \rangle| < \eta$ for $i < j$.

Since $f_n \xrightarrow{w^*} f$, we can find a subsequence $1 < k_1 < k_2 < \dots < k_n < \dots$ such that $|\langle x_1, f_m - f \rangle| < \eta$ for $m > k_1$, $|\langle x_{k_1}, f_m - f \rangle| < \eta$ for $m > k_2$, and so on.

So, the subsequence $1 < k_1 < k_2 < \dots < k_n < \dots$ satisfies $|\langle x_{k_i}, f_{k_j} - f \rangle| < \eta$ for $i < j$.

Without loss of generality, we still use \mathbb{N} to denote the subsequence $1 < k_1 < k_2 < \dots < k_n < \dots$. So we have $|\langle x_i, f_j - f \rangle| < \eta$ for $i < j$.

It follows that $|\langle x_i, f_j \rangle| \leq |\langle x_i, f_j - f \rangle| + |\langle x_i, f \rangle| < 2\eta = \varepsilon$ if i and j large enough and $i < j$.

By deleting some terms in the front if necessary, we have $|\langle x_j, f_i \rangle| < \varepsilon$ for $i \neq j$.

Finally, $\|f_i - f_j\| \geq |\langle x_j, f_i - f_j \rangle| = |1 - \langle x_j, f_i \rangle| \geq 1 - |\langle x_j, f_i \rangle| > 1 - \varepsilon$ for $i \neq j$. \square

Theorem 2.19. *Let X be a Banach space such that $B(X^*)$ is weak* sequentially compact. If $\text{UKK}^*((\frac{1}{\mu(X^*)})^-) > 1 - \frac{1}{\mu(X^*)}$ for X^* , then X has weak normal structure.*

Proof. Suppose that X fails to have weak normal structure. From Lemma 2.18, for any $\eta > 0$, let the sequence $\{x_1, x_2, x_3, \dots, x_n, \dots\} \subseteq S(X)$ with $x_n \xrightarrow{w} 0$ and the sequence $\{f_1, f_2, \dots, f_n, \dots\} \subseteq S(X^*)$ with $f_n \xrightarrow{w^*} f$ satisfy four conditions there.

It is easy to see that from the definition of $\mu(X^*)$, we can find a subsequence $1 < k_1 < k_2 < \dots < k_n < \dots$ such that $1 - \eta < \|f_{k_i} - f_{k_j}\| < \mu(X^*) + \eta$ for $i \neq j$.

Without loss of generality, we still use \mathbb{N} to denote the subsequence $1 < k_1 < k_2 < \dots < k_n < \dots$. So we have $1 - \eta < \|f_i - f_j\| < \mu(X^*) + \eta$ and $|\langle x_j, f_i \rangle| < \eta$ for all $i \neq j$.

Consider the sequence $\{g_1 = f_1 - f_2, g_2 = f_1 - f_3, \dots, g_n = f_1 - f_{n+1}, \dots\} \subseteq B_{\mu(X^*) + \eta}(0)$ for X^* . Then $f_n - f \xrightarrow{w^*} 0$. Moreover, for each i , $|\langle x_i, f \rangle| = \lim_{n \rightarrow \infty} |\langle x_i, f_n \rangle| < \eta$. So, $\|f_n - f\| \geq |\langle x_n, f_n - f \rangle| > 1 - \eta$ for each n . Note that $1 - \eta \leq \|g_i - g_j\| = \|f_{j+1} - f_{i+1}\| \leq \mu(X^*) + \eta$ for $i \neq j$, and $g_n \xrightarrow{w^*} f_1 - f$.

Let $z_i = \frac{g_i}{\mu(X^*)}$ for all $i \in \mathbb{N}$. Then

$$\frac{1}{\mu(X^*)} - \frac{\eta}{\mu(X^*)} \leq \|z_i - z_j\| = \frac{\|g_i - g_j\|}{\mu(X^*)} \leq 1 + \frac{\eta}{\mu(X^*)} \quad \text{and} \quad z_n \xrightarrow{w^*} \frac{f_1 - f}{\mu(X^*)}.$$

From the definition of modulus of UKK^* for X^* , we have $\text{UKK}^*(\frac{1}{\mu(X^*)} - \frac{\eta}{\mu(X^*)}) < 1 - \frac{1-\eta}{\mu(X^*)}$. Since η can be arbitrarily small,

$$\text{UKK}^*\left(\left(\frac{1}{\mu(X^*)}\right)^-\right) < 1 - \frac{1}{\mu(X^*)}.$$

This completes the proof. \square

Let $\mu(X^*) = 1$ in Theorem 2.19. Then we obtain the following result.

Theorem 2.20. *Let X be a Banach space such that $B(X^*)$ is weak* sequentially compact. If $\text{UKK}^*(1^-) > 0$ for X^* and $\mu(X^*) = 1$, then X has weak normal structure.*

We now consider the ultraproduct spaces. For an ultrafilter \mathcal{U} on \mathbb{N} , the set of natural numbers, we use $X_{\mathcal{U}}$ to denote the ultraproduct. For more details, see [19].

If X is super-reflexive, then $\tilde{X}^* = (\tilde{X})^*$ and X has uniform normal structure if and only if \tilde{X} has normal structure [19]. Since X can be embedded into \tilde{X} , it is easy to see that $\text{NUC}(\varepsilon)$ of X is greater than or equal to $\text{NUC}(\varepsilon)$ of \tilde{X} . By using these facts, we can prove the following result about uniform normal structure.

Theorem 2.21. *Let X be a super-reflexive Banach space with $\text{NUC}(1^-) > 0$ for \tilde{X} . Then X has uniform normal structure.*

Proof. If X is a super-reflexive Banach space but fails to have uniform normal structure, then \tilde{X} fails to have normal structure. From Theorem 2.14, we have $\text{NUC}(1^-) > 0$ for \tilde{X} . \square

Acknowledgements

The authors are very grateful to the referees for several helpful suggestions and improvements. The first author was supported by the Centre of Excellence in Mathematics and the Commission on Higher Education, Thailand.

References

- [1] M.S. Brodskii, D.P. Mil'man, On the center of a convex set, Dokl. Akad. Nauk SSSR (NS) 59 (1948) 837–840 (in Russian).
- [2] W.A. Kirk, A fixed point theorem for mappings which do not increase distances, Amer. Math. Monthly 72 (1965) 1004–1006.
- [3] R. Huff, Banach spaces which are nearly uniformly convex, Rocky Mountain J. Math. 10 (4) (1980) 743–749.
- [4] D. van Dulst, B. Sims, Fixed points of nonexpansive mappings and Chebyshev centers in Banach spaces with norms of type (KK), in: Banach Space Theory and its Applications, Bucharest, 1981, in: Lecture Notes in Math., vol. 991, Springer, Berlin, New York, 1983, pp. 35–43.
- [5] P.N. Dowling, B. Randrianantoanina, B. Turett, The fixed point property via dual space properties, J. Funct. Anal. 255 (3) (2008) 768–775.
- [6] J.M. Ayerbe Toledano, T. Domínguez Benavides, G. López Acedo, Measures of Noncompactness in Metric Fixed Point Theory, in: Operator Theory: Advances and Applications, vol. 99, Birkhäuser Verlag, Basel, 1997.
- [7] J. Banaś, On modulus of noncompact convexity and its properties, Canad. Math. Bull. 30 (2) (1987) 186–192.
- [8] J. Banaś, L. Olszowy, K. Sadarangani, Moduli of near convexity of the Baernstein space, Proc. Amer. Math. Soc. 123 (12) (1995) 3693–3699.
- [9] T. Domínguez Benavides, G. López Acedo, Lower bounds for normal structure coefficients, Proc. Roy. Soc. Edinburgh Sect. A 121 (3–4) (1992) 245–252.
- [10] K. Goebel, T. Sekowski, The modulus of noncompact convexity, Ann. Univ. Mariae Curie-Skłodowska Sect. A 38 (1984) 41–48. 1986.
- [11] H. Knaust, E. Odell, Th. Schlumprecht, On asymptotic structure, the Szlenk index and UKK properties in Banach spaces, Positivity 3 (2) (1999) 173–199.
- [12] B. Sims, M.A. Smyth, On some Banach space properties sufficient for weak normal structure and their permanence properties, Trans. Amer. Math. Soc. 351 (2) (1999) 497–513.
- [13] J.R. Partington, On nearly uniformly convex Banach spaces, Math. Proc. Cambridge Philos. Soc. 93 (1) (1983) 127–129.
- [14] D. van Dulst, Some more Banach spaces with normal structure, J. Math. Anal. Appl. 104 (1) (1984) 285–292.
- [15] C.A. Kottman, Packing and reflexivity in Banach spaces, Trans. Amer. Math. Soc. 150 (1970) 565–576.
- [16] R.C. James, Weakly compact sets, Trans. Amer. Math. Soc. 113 (1964) 129–140.
- [17] S. Saejung, J. Gao, On semi-uniform Kadec–Klee Banach spaces, Abstr. Appl. Anal. 2010, 12 pages, Article ID 652521, doi:10.1155/2010/652521.
- [18] S. Saejung, Convexity conditions and normal structure of Banach spaces, J. Math. Anal. Appl. 344 (2) (2008) 851–856.
- [19] B. Sims, Ultra-Techniques in Banach Space Theory, in: Queen's Papers in Pure and Applied Mathematics, vol. 60, Queen's University, Kingston, ON, 1982.